

Infrared Abelian Dominance and Dual Higgs Mechanism in MA Gauge

H. Suganuma^{a*}, K. Amemiya^a and H. Ichie^a

^aResearch Center for Nuclear Physics (RCNP), Osaka University, Ibaraki, Osaka 567-0047, Japan

We study infrared abelian dominance and the dual Higgs mechanism in the maximally abelian (MA) gauge using the lattice QCD Monte Carlo simulation. In the MA gauge, the off-diagonal gluon phase tends to be random, and the off-diagonal gluon $A_\mu^\pm$ acquires the effective mass as $M_{\text{off}} \simeq 1.2$ GeV. From the monopole current in the MA gauge, we extract the dual gluon field B_μ and estimate the dual gluon mass as $m_B \simeq 0.5$ GeV. The QCD-monopole structure is also investigated in terms of off-diagonal gluons. From the lattice QCD in the MA gauge, the dual Ginzburg-Landau (DGL) theory can be constructed as a realistic infrared effective theory based on QCD.

1. Strong Randomness of Off-diagonal Gluon Phase in MA gauge

We find *strong randomness of the off-diagonal gluon phase* χ_μ in the MA gauge using the SU(2) lattice QCD [1,2]. In fact, we find very small correlation between neighboring phases, $\chi_\mu(s)$ and $\chi_\mu(s+\hat{\nu})$, on the lattice in the MA and U(1)₃ Landau gauge. This tendency seems natural because of the following reasons. Off-diagonal gluon phase χ_μ is not constrained by the MA gauge-fixing condition at all, and the constraint from the QCD action is also small because of strong suppression of the off-diagonal gluon amplitude $|A_\mu^\pm|$ in the MA gauge. Within the random phase-variable approximation, *perfect abelian dominance for the string tension* can be clearly demonstrated [1,2].

2. Propagator and Effective Mass of Off-diagonal Gluon in MA gauge

Using the SU(2) lattice QCD with $12^3 \times 24$, 16^4 and 20^4 , we study the off-diagonal gluon propagator $G_{\mu\mu}^{\text{off}}(r) \equiv \langle A_\mu^+(x) A_\mu^-(y) \rangle$ as the function of $r \equiv \sqrt{(x-y)^2}$ in the MA gauge with U(1)₃ Landau gauge fixing [2,3]. From the slope of the logarithmic plot of $r^{3/2} G_{\mu\mu}^{\text{off}}(r)$ in Fig.1(a), we get the off-diagonal gluon mass as $M_{\text{off}} \simeq 1.2$ GeV in the MA gauge [2,3] as shown in Fig.1(b). Thus, in the MA gauge, the off-diagonal gluon cannot carry the long-range interaction such as the con-

finement force. This is the essence of infrared abelian dominance. In fact, QCD in the MA gauge behaves as a strong-coupling compact QED at the larger scale than $M_{\text{off}}^{-1} \simeq 0.2$ fm [2,3].




Figure 1. (a) The logarithmic plot of $r^{3/2} G_{\mu\mu}^{\text{off}}(r)$, $r^{3/2} G_{\mu\mu}^{\text{Abel}}(r)$. The slope corresponds to the effective mass. (b) M_{off} v.s. inverse lattice volume.

3. Dual Gluon Mass in MA gauge

Using the SU(2) lattice QCD with 20^4 and 24^4 , we extract the dual gluon field B_μ from the monopole current k_μ appearing in the MA gauge, and investigate the dual gluon properties [2].

*e-mail: suganuma@rcnp.osaka-u.ac.jp

(1) From the dual Wilson loop, we obtain the inter-monopole potential in the monopole part.
 (2) The dual gluon propagator $\langle B_\mu(x)B_\mu(y) \rangle$ is measured in the dual Landau gauge $\partial_\mu B_\mu = 0$.
 As the evidence of the dual Higgs mechanism by color-magnetic monopole condensation, the dual gluon mass is estimated as $m_B \simeq 0.5$ GeV from the lattice data in $0.4 \text{ fm} < r < 1.5 \text{ fm}$ [2]. Here, the dual gluon mass $m_B \simeq 0.5$ GeV provides the key parameter of the DGL theory.




Figure 2. (a) The inter-monopole potential. (b) $\ln(r_E^{3/2} \langle B_\mu(x)B_\mu(y) \rangle)$ v.s. 4 dim. distance r_E .

4. Structure of QCD-Monopoles in terms of Off-diagonal Gluons

We investigate the QCD-monopole structure in the MA gauge in the lattice QCD [2].

(1) Around the QCD-monopole, both the abelian action density S_{Abel} and the off-diagonal gluon contribution $S_{\text{off}} \equiv S_{\text{QCD}} - S_{\text{Abel}}$ are largely fluctuated, and their cancellation keeps the total QCD action density S_{QCD} small [2].

(2) The QCD-monopole has an intrinsic structure relating to a large amount of off-diagonal gluons around its center, similar to the 't Hooft-Polyakov monopole [2]. At a large scale where this structure becomes invisible, QCD-monopoles can be regarded as point-like Dirac monopoles.

(3) From the concentration of off-diagonal gluons around QCD-monopoles in the MA gauge, we can naturally understand the *local correlation between monopoles and instantons*. In fact, instantons tend to appear around the monopole world-line in the MA gauge, because instantons need full SU(2) gluon components for existence [2,4].

5. Construction of the DGL theory from the lattice QCD in MA gauge

In the MA gauge, the off-diagonal gluon contribution can be neglected and monopole condensa-


tion occurs at the infrared scale of QCD. Therefore, the QCD vacuum in the MA gauge can be regarded as the dual superconductor described by the DGL theory, and quark confinement can be understood with the dual Meissner effect. The DGL theory can describe not only quark confinement [5] but also dynamical chiral-symmetry breaking (D χ SB) by solving the Schwinger-Dyson equation for the quark field [5,6]. Monopole dominance for D χ SB is also found from the analysis of the effective potential [7]. Using the DGL theory, we have studied the QCD phase transition (deconfinement [5,8] and chiral restoration [6]) at finite temperature, the QGP creation process in ultra-relativistic heavy-ion collisions [5,8], hadronization in the early universe [9], and glueball properties [10] systematically.

REFERENCES

1. H. Ichie and H. Suganuma, *Nucl. Phys.* **B548** (1999) 365; *Phys. Rev.* **D60** (1999) 77501.
2. H. Suganuma, H. Ichie, A. Tananka and K. Amemiya, *Prog.Theor.Phys.Suppl.* **131** (1998) 559; H. Suganuma, K. Amemiya, H. Ichie, A. Tananka, *Nucl. Phys.* **A** (1999).
3. K. Amemiya, H. Suganuma, this Proceedings.
4. M. Fukushima, H. Suganuma and H. Toki, *Phys. Rev.* **D60** (1999) 94504; M. Fukushima, S. Sasaki, H. Suganuma, A. Tanaka, H. Toki, D. Diakonov, *Phys. Lett.* **B399** (1997) 141.
5. H. Suganuma, S. Sasaki and H. Toki, *Nucl. Phys.* **B435** (1995) 207 and its references ; H. Suganuma, S. Sasaki, H. Toki and H. Ichie, *Prog. Theor. Phys. Suppl.* **120** (1995) 57.
6. S.Sasaki, H.Suganuma, H.Toki, *Prog. Theor. Phys.* **94** (1995) 373; *Phys. Lett.* **B387** (1996) 145; *Prog.Theor.Phys.Suppl.* **129** (1998) 227.
7. S. Umisedo, H. Suganuma and H. Toki, *Phys. Rev.* **D57** (1998) 1605.
8. H. Ichie, H. Suganuma and H. Toki, *Phys. Rev.* **D52** (1995) 2944; **D54** (1996) 3382.
9. H. Monden, H. Ichie, H. Suganuma and H. Toki, *Phys. Rev.* **C57** (1998) 2564.
10. Y. Koma, H. Suganuma, H. Toki, *Phys. Rev.* **D60** (1999) 74024; Y. Koma *et al.*, *NPA*.

From Lattice QCD to Dual Ginzburg-Landau Theory

— Infrared Effective Theory directly based on QCD —



Dual Gauge Symmetry is spontaneously broken instead of Gauge Symmetry

Figure 3. Construction of the dual Ginzburg-Landau (DGL) theory from the lattice QCD in